

THE AMOUNT OF SOME MINERAL SUBSTANCES IN COWS' BLOOD SERUM FED WITH CORN SILAGE**

J. Stojkovic^{1*}, B. Jasovic¹, D. Randjelovic¹

¹ Faculty of Agriculture Pristina, Serbia

* Corresponding author: jovanips@bankerinter.net

** Communications

Abstract: The amount of calcium, phosphorus and magnesium in milking cows' blood was tested and it is one of indicators that cows were fed with these substances. The average amount of calcium in tested cows' blood is 11.58mg/100ml serum, the average amount of phosphorus is 5,14mg/100ml and the average amount of magnesium is 3,03/100mg .

Key words: cow, calcium, phosphorus and magnesium.

Introduction

The amount of these mineral substances in large content of fodder all over our country is not tested enough. Last research results of mineral substances in the fodder show the low level of phosphorus that appears more often than calcium does .This manifestation appears more often during the drought and during some periods that occurs very often at Kosovo and Metohija and where the tests were performed. It should be mentioned the fact that the soil at Kosovo and Metohija contains a low level of physiologically active phosphorus (P₂O₅), as *Miric* (1989) established and the low level of this element in soil has an adverse affect on the level of this element in plants. It should be mentioned Stojkovic's observation that lots of plants from this area contain the low level of phosphorus. Examinations of the level of calcium, phosphorus and magnesium in cows' blood serum which are fed by different fodder as a source of these elements and standard fodder with mineral would certainly contribute to better knowingness of general status and dynamics circulation of these elements and their relations in cows' blood serum. They assumed that their level in blood serum is one of the indicators that cows are fed with these elements. Contribution is certainly greater since the cows with different lactation were examined.

Material and methods

The experiments were carried out on Simmental cows at the farm. It lasted 45 days during which the blood samples were taken for the mineral analyses. There were 20 cows and five of them were at the beginning of the lactation period with average 25 kg of milk per day (first group). In the second group there were five cows in the middle of lactation period with average 22,5kg of milk per day. In the third group there were five cows at the end of lactation period with average 12,6 kg of milk per day and in fifth group were pregnant cows.

They were fed with 4.0 kg of alfalfa hay , 25.0 kg of maize silage, while the fodder for the milking cows was distributed by groups : cows at the beginning of lactation 5kg , cows in the middle of lactation 4 kg, and those at the end of lactation and pregnant cows 3 kg. Structure and nutrition value of cows' servings are showed in the table 1 and chemical composition, nutritious value and mineral composition in table 2 .

Table 1. Composition and nutritious value of serving.

Group	Food	kg	S.M.(kg)	H.J.	S.P.(g)	Ca (g)	P(g)	Mg(g)
I	<i>Hay lucerne</i>	4,0	3,64	1,82	337	41,0	13,8	2,90
	<i>Maize silage</i>	25,0	7,50	7,20	520	127,0	45,3	11,8
	<i>Fodder</i>	5,0	4,55	5,60	587	75,0	51,9	4,60
	<i>Total</i>		15,69	14,62	1444	243,0	111	19,0
II	<i>Hay lucerne</i>	4,0	3,64	1,82	335	41,0	13,8	2,95
	<i>Maize silage</i>	25,0	7,50	7,0	495	130,0	45,3	11,89
	<i>Fodder</i>	4,0	3,50	4,85	470	61,2	37,6	3,80
	<i>Total</i>		14,64	13,66	1300	232,2	96,7	18,64
III and IV	<i>Ha lucerne</i>	4,0	3,64	1,82	337	41,0	13,8	2,90
	<i>Maize silage</i>	25,0	7,50	7,0	495	130,0	45,3	11,70
	<i>Fodder</i>	3,0	2,66	4,10	380	52,2	28,9	3,15
	<i>Total</i>		13,80	12,92	1212	224,2	88,0	17,75

The level of calcium, phosphorus and magnesium in cows' blood and food they were fed with is tested. Cows' blood was taken v. Jugularis and the serum was isolated and tested to these metals. Calcium in the samples of fodder was tested by permanganate method. Serum samples were tested by atomic -absorbing methods, the hollow cathode lamp HCL at 442.7 nanometre wave lenght. Phosphorus in the fodder samples was tested by classical gravimetical methods, measured as $Mg_2P_2O_7$, and serum samples by colourmertical methods on photocolourmetre. Magnesium in the fodder samples was tested by gravimetical method as MgO , and serum on atomic absorber.

Table 2. Chemical composition and nutritious value of used foodstuff.

Food	Hay lucerne	Maize silage	Fodder
Moisture%	9,45	69,00	11,30
Ash %	6,30	3,59	6,50
Proteins %	16,80	4,22	12,50
Cellulose%	31,00	7,98	5,87
Fat %	3,80	0,34	3,50
BEM %	33,85	16,87	60,70
Ca (g)	13,74	5,20	13,90
P (g)	4,50	1,80	6,83
Mg (g)	0,95	2,0	1,10
S.P. (g)	1,10	1,98	11, 50
H.J	0,45	0,28	1,10

Statistical analysis of data is done by Statistica programme version 6, StatSoft. Inc. (2003) .

Research results and Discussion

The amounts of calcium, phosphorus and magnesium, measured in the experiments, are in table 3 and 4.

The average amount of calcium in cows' blood serum in the experiment is 11.83 mg per 100mg milk (table 4). That amount was within the normal limits. The amount of calcium in fodder was sufficient to provide enough calcium in the blood serum. (224-243 g).

The average amount of phosphorus for all groups in the experiment is 5.14 mg per 100mg milk. This result is on the lower level within the normal scale amount based on literature data (4,5-6,5 mg/100 ml).Observing apart

the amounts of the content it can be seen that 9 out of 20 cows had amount of phosphorus under the lower limit of normal content. So it can be concluded that there was not enough phosphorus in the foodstuff. Since it was established that there was enough amount of phosphorus in the fodder, it is possible that other factors influenced to the low amount in the blood serum among which it should be mentioned first of all the level of usage out of some kind of fodder.

Based on the mineral elements' analysis, it is concluded that amount of calcium and phosphorus are directly connected to cows' lactation. The group of cows with average daily lactation of 25.0 kg had the lowest amount of calcium and the pregnant cows had the highest amount. The proportion was 10.52:12,40 mg/100 ml serum (tabela 3).

The amount of magnesium was within the normal limits (3.15mg/100) and there were slight changes among groups and among cows. There were no individual changes in the amount of magnesium nor there were changes within the group of cows with different lactation. But groups' average amounts indicated that magnesium is not in the same relations to lactation as calcium and phosphorus are. It means that the cows with highest lactation had the highest amount of magnesium and vice versa. The difference between the lowest and the highest amount was 9%, in the favor of the highest amount.

Table 3. Content of calcium, phosphorus and magnezium in blood serum of investigated cows (mg/100ml)

Group	Mineral matters	X
I	Ca	10,52
	P	4,08
	Mg	3,20
II	Ca	11,65
	P	4,81
	Mg	3,18
III	Ca	10,85
	P	5,60
	Mg	3,12
IV	Ca	12,40
	P	6,10
	Mg	3,10

Judging by data from the book *Underwood* (1972, 1976), *Mirića* (1990), this amount was 8.66% lower from the average amount of calcium in the

previous experiments *Stojković* (2001, 2006). Based on same data the amount of calcium in the blood serum of milking cows can be considerably higher, *Vijchulata i sar.* (1983) which was the case when high amounts of calcium were added to their servings. There are also cases with extremely low amounts of calcium. As *Lalov and assos.* (1976) mentioned in his experiments the amount of calcium was just 4.47 mg in 100 ml with the very high amount of phosphorus (15,67 mg/100) at the same time.

Table 4. The content of calcium, phosphorus and magnesium in the cows' blood serum, average amount for all groups

Element	X	S	CV	IV
Ca	11,35	2,71	11,30	8,97 – 13,17
P	5,14	2,50	18,90	3,68 – 7,12
Mg	3,15	2,10	8,90	2,10-3,62

The amount of phosphorus changed in the relation to cows' lactation. The group of cows with average daily lactation of 25.00 kg had the lowest amount of phosphorus in the blood serum which was 4.08mg/100ml serum (table 3). These amounts were under the lower limits of normal amounts and also insufficient amounts for the organism. The amount of phosphorus was increasing in the group of cows with lower lactation. In the group of cows with average daily lactation of 22.50 kg the amount of phosphorus was 4.81 mg/100 ml and that was above lower level of the normal amounts. This amount was even more increased (5.60 mg in 100 ml) with the group of cows with average daily lactation of 12.6kg and even more (6.10 mg/100 ml) (table 3) with the group of pregnant cows.

In the experiment with milking cows *Mirić and assos.* (1990) established that the amount of phosphorus in the blood serum is in correlation with lactation which *Stojković's* (2001, 2006) results confirmed. The amount of phosphorus in *Stojković's* previous testing was 5% lower than in this. The fact that amount of phosphorus in the blood serum can be above the average (4.5-6.5 mg/100 ml) confirms the experiments of *Underwoodu* (1972, 1976) and *Heraka and assos.* (1976) where the pregnant heifers had from 6.92 to 7.8 mg% of phosphorus. The amount of phosphorus was even greater (9.94-10.40 mg/100 ml) in the experiments of *Seidela and assos.* (1970).

Underwoodu (1976) and Obračeviću (1973) explained that there were slight changes in the amount of magnesium because magnesium doesn't decrease in the soft tissues even when skeleton lost 30% of magnesium. The results of experiments showed certain correlation between the amount of calcium and phosphorus and lactation. The cows with high lactation had lower amounts of these elements and vice versa. This fact was established even when these elements were not examined or showed a deficit. This manifestation was evident in the variations in the amount of phosphorus and little less evident in the amount of calcium while the amount of magnesium behaved differently.

Conclusion

The levels of calcium, magnesium and phosphorus in the blood serum were tested, as one of the indicators that the cows are fed with these elements.

The results of the experiments are :

1. The average amount of calcium in the blood serum is 11.35mg in 100 ml of serum, of phosphorus is 5.14 mg/100 ml and magnesium 3.15 mg/100 ml. The amounts of calcium were on the upper level of normal amounts gotten from books while the amount of phosphorus was almost deficient.
2. The amounts of calcium and phosphorus were decreasing during the period of high lactation while the cows with low lactation had these amounts increased. Calcium varies from 10.52 to 12.40 and phosphorus varies from 4.08 to 6.10 mg in 100 ml of serum.
3. The amounts of magnesium didn't change much in relations to cows' lactation but magnesium behaved differently from calcium and phosphorus.

SADRŽAJ NEKIH MINERALNIH MATERIJA U KRVNOM SERUMU KRAVA HRANJENIH KUKURUZNOM SILAŽOM

J. Stojković, B. Jašović, D. Randjelovic

Rezime

Ispitivan je sadržaj kalcijuma, fosfora i magnezijuma u krvnom serumu muznih krava, kao jednog od indikatora obezbeđenosti životinja ovim materijama. Prosečan sadržaj kalcijuma u krvnom serumu ispitivanih krava iznosio je 11,58 mg/100 ml seruma, fosfora 5,14 mg/100 ml i magnezijuma 3,03 mg u 100 ml seruma. Krave su hranjene kukuruznom silažom, senom lucerke i krmnom smešom za krave muzare.

Ključne reči: krava, kalcijum, fosfor, magnezijum

References

- HERAK M., HERAK M., BELIĆ J., VRBANAC I., ČAUŠEVIĆ N., KOVAČEVIĆ V., ANDRAK T. (1976): Odnos hormonalnog statusa organizma i promjene razine bakra, kalcija, fosfora i magnezija u krvnom serumu goveda i svinja. Zbornik radova Poljoprivrednog fakulteta u Sarajevu, 27.
- LALOV H., CVETKOV A., ANTONOV S., DIMITROV A., GORANOV H., MALČEVSKI M. (1976): Proučavane smuščenijata v kalcijevofosfornija obmen pri intenzivno ugojavane na mladi goveda. Veterinarnomed. Nauki, 13, 1, 8-22.
- MIRIĆ M. (1989): Krmna baza Kosova. Monografija. Priština.
- MIRIĆ M., JOKIĆ A., STOJKOVIĆ J. (1990): Uticaj kukuruzne silaže u obroku krava muzara na sadržaj kalcijuma, fosfora i magnezijuma u krvnom serumu. Stočarstvo 44:1990 (3-4) 101-107.
- OBRAČEVIĆ Č. (1973): Osnovi ishrane domaćih životinja. Minerva, Subotica – Beograd.
- SEIDEL H., SCHRUES J. (1970): Das Verhalten der Natrium, Kalium, Calcium, Magnesium und Phosphor konzentration im Serum des graviden und laktierenden Rindes. Arch. Exp. Veterinarmed, 24, 873-882.
- STOJKOVIĆ J., RAJIĆ I., RADOVANOVIĆ T. (1996): Pregled i ocena stočne hrane. Monografija. Priština.
- STOJKOVIĆ J. (2001): Sadržaj kalcijuma, fosfora i magnezijuma u krvnom serumu krava različite mlečnosti. Poljoprivreda i šumarstvo, Vol 47. (3-4), 113-120. Podgorica.
- STOJKOVIĆ J. (2006): Sadržaj kalcijuma i fosfora u krvnom serumu krava hranjenih zelenim sirkom. Biotehnologija u stočarstvu, vol. 22, 651-658. Beograd.

STOJKOVIĆ J. (2006): Mineralne materije u ishrani domaćih životinja. Monografija. Niš.

Statistica, verzion 6, StotSoft. Inc (2003).

UNDERWOOD F.J. (1972): Trace elements in Human and Animal Nutrition. Academic Pres Inc. New York and London.

UNDERWOOD F.J. (1976): Trace mineral imbalances in farm animals and their study and diagnosis with isotopic tracers Atom Energy Rev., 14.

VIJCHULATA P., SRISAK T., PRUKSASRI P. (1983): Effects of mineral supplements on growth and profile of minerals in growing cattle grazed in paragrass pasture. New. Strateg. Improvv. Animal Prod. Hum. Welfare. Proc. World Conf. Anim. Prod. Tokyo, 14-19.