

BIOLOGICAL AND TECHNOLOGICAL BACKGROUND OF ESTRUS SYNCHRONIZATION AND FIXED-TIME OVULATION INDUCTION IN THE PIG

K.-P. Brüssow¹, M. Wähler²

¹Leibniz Institute for Farm Animal Biology (FBN), Dummerstorf, Germany

²Anhalt University of Applied Sciences, Bernburg, Germany

Corresponding author: bruessow@fbn-dummerstorf.de

Invited paper

Abstract: A technology that allows for manipulating of estrus and ovulation, and would then also allow for fixed-time insemination, can be of great benefit for swine farms that operate using sow batch management. Such technology at least in part, saves labor and permits the production of large batches of evenly developed pigs. Thanks to the current knowledge on endocrine regulation of follicle development and ovulation, and the availability of numerous reproductively active substances such a technology is now available. This 'biotechnology of reproduction' will be reviewed. It covers procedures for synchronizing estrus based on the use of altrenogest in gilts and of batch-wise weaning in sows, for stimulating follicle development using eCG and for inducing of ovulation using hCG or LH as well as GnRH analogues. While the procedures for estrus synchronization stand alone, other procedures require additional treatments. If fixed-time insemination is the goal, estrus needs to be synchronized and follicular development and ovulation induced by the use of GnRH analogues and hCG with ovulation occurring within 36-42 hrs. It is a general recommendation to inseminate those animals twice, i.e. 24 and 40 hrs after ovulation induction. However, the aforementioned technology requires healthy animals and a solid management and cannot be used to compensate for poor management.

Key words: estrus synchronization

Introduction

More than half a century ago research efforts were made to synchronize the estrous cycle and ovulation in pigs with the ultimate goal of fixed-time artificial insemination (AI). *Tanabe et al. (1949)* were the first to treat pigs with equine chorionic gonadotrophin (eCG) and sheep pituitary extracts in order to stimulate follicle development and ovulation. Further studies, which were milestones in the

development of procedures to manipulate the estrus cycle in pigs, were those conducted by *Polge and Day (1969)* using methallibure to restrain follicular growth followed by treatment with eCG to stimulate follicle development and human chorionic gonadotrophin (hCG) for ovulation induction, and by *Hunter (1967, 1974)* on follicle and oocyte development. Results of these early studies facilitated Eastern-European, particularly East-German research efforts into those procedures starting in 1970 at a time when the farm inventories were growing and there was an increasing need for this kind of biotechnology for management purposes. The general approach was to manipulate female reproductive functions, such as follicular development as well as ovulation and parturition. However, the goal was to mimic what occurs physiologically in the female pig. The ultimate goal was to synchronize all reproductive processes with the advantage of periodic and batch-wise AI, parturition as well as weaning, thereby enabling the practice of all-in-all-out and to produce large groups of pig in the same reproductive state with the same health and immunization status. Indicative of the success made over the years in reproductive biotechnology in 1990, controlled reproduction was used on 86 % of the 1.1 million breeding sows and gilts in East-Germany. It has been learned that research into and the practical use of biotechnological procedures requires understanding of reproductive processes and the availability of appropriate substances (i.e. hormones or hormonally active substances). While those substances are available in general to manipulate almost all key reproductive processes in the female pig, they are not equally available since their use in practice requires national approval. As compared to Europe, hormones or hormonally active substances are still rarely used in the North and South American as well as the Asian swine industry. However, during the last decade due to increased costs for labor, feed and energy resulted in new interest and demand in using those substances that allow for manipulating reproductive processes in the female pig.

Follicle development and ovulation

Follicular development in gilts and sows has been reviewed in detail elsewhere (*Prunier and Quesnel, 2000; Schwartz et al., 2008*). In gilts the follicle cohort destined to ovulate is stimulated by increased postovulatory FSH, but under the influence of the high luteal progesterone (P4) concentration during dioestrus follicles do not grow to ovulatory size. Only after luteolysis when P4 is low and thus does not negatively feedback on gonadotrophin synthesis, follicles finally grow from 4 mm to ovulatory size within 4-6 days. However, there is a well-defined balance between stimulatory (e.g. LH and FSH) and inhibitory (e.g. P4 and inhibin) factors that favors preovulatory follicle development. As has been shown after gonadotrophin deprivation, FSH is necessary to support follicle development

beyond 2-3 mm and LH beyond 4 mm (*Driancourt et al., 1995; Brüßow et al., 2001*). Once the follicles reach a larger preovulatory stage FSH declines (*Guthrie and Bolt, 1990*) and LH pulse secretion changes from luteal (high amplitude - low frequency) to follicular patterns (low amplitude - high frequency). Final stage of follicular development is associated with decreased FSH and increased LH receptor expression along with increased production of estradiol, as well as increased fluid accumulation within the follicular cavity. Increased preovulatory estradiol finally elicits the GnRH-mediated LH release from the pituitary, thereby initiating ovulation and the release of a mature oocyte.

Sows commonly have a lactational anestrus. During lactation and before weaning, follicles exhibit a wave-like pattern of growth, and emerge from a cohort of 20-30 follicles of 2 mm that grow to not larger than 5 mm (*Lucy et al., 2001*). This is because suckling inhibits secretion of GnRH and subsequently LH due to a concerted action of prolactin, oxytocin and endogenous opioid peptides that prevent final growth of these follicles to reach ovulatory size before weaning (*Varley and Foxcroft, 1990*). Once the piglets are weaned, follicles start to grow to 7-8 mm before ovulation. Initially FSH increases and then decreases at weaning. Basal LH and LH pulse frequency increases; these are key regulators of post-weaning follicle development which affect the weaning to estrus interval (*van de Brand et al., 2000*). Furthermore, adrenal hormones and nutritional mediators such as glucose, insulin and free fatty acids, as well as endogenous opioids and leptin have been shown to have an influence on post-weaning follicle development and ovulation. In conclusion, follicle development including the final ovulatory follicle growth and maturation, as well as ovulation itself, requires fine-tuned hormonal events. Understanding these mechanisms is necessary in any attempt to manipulate the reproductive process in female swine.

Manipulation of follicle development and ovulation

Synchronization of estrus. In a batch farrowing system, estrus in sows is naturally synchronized by weaning. Estrous cycles of gilts need to be synchronized when they are introduced into such systems. For this purpose, protocols based on either suppression of follicle development and/or mimicking the luteal phase are available (reviewed by *Estill, 1999*). Progesterone and its synthetic derivatives did not prove fully effective for estrus synchronization. In contrast, non-steroidal substances such as methallibure (Aimax, ICI 33828, Suisynchron®) have been shown to yield good results, but were banned by US and European countries before being commercially available for the swine industry because of their teratogenic effects. In East-Germany, zinc-methallibure (Siusynchron®) was successfully used in hundreds of thousands of gilts between 1973 and 1989, but was then also banned and replaced by allyltrenbolone (*Hühn et al., 1996*). At present, allyltrenbolone (altrenogest, as Regumate® in Europe and Matrix® in North America) is the only

licensed substance with progestagene type effects to be used in female pigs in Europe and in North America. If it is given orally and fed in a dose of 15 to 20 mg/day/gilt over a period of 14 to 18 days, it has been proven effective at suppressing follicle development. Recommendations on the daily dose of altrenogest and the duration of administration differ between countries. Production sites in France and also some in Germany prefer feeding 20 mg/day/gilt for 18 days (*Martinat-Botte et al., 1990*). Other sites in Germany use feeding altrenogest in doses of 16-20 mg/day/gilt over 15 days and achieved good fertility results (*Hühn et al., 1996*). For North America, the general recommendation is 15 mg/day/gilt for 14 days. Feeding of altrenogest for a shorter period is possible, if the stage of estrous cycle is known at first administration (*Kirkwood, 1999*). Gilts usually show estrus within 5 to 7 days after altrenogest withdrawal (*Martinat-Botte et al., 1990; Hühn et al., 1996*). Other approaches have been to include progesterone or GnRH agonists in a slow release device during a defined time, in conjunction with hCG or estrogen to extend the lifespan of the corpora lutea, and the 'breed-and-abort' procedure using PGF2 α . However, all of the aforementioned procedures except those based on the use of altrenogest are not currently used in the swine industry.

Stimulation of follicular development. Though synchronization of estrus is possible, with weaning group of sows or treatment with altrenogest in gilts, the onsets of estrus can however still spread over a week. This is due, at least in part, to insufficient follicle development. Gonadotrophins may then be used after weaning or after altrenogest in order to stimulate this follicle development and to achieve a better synchronization effect. Equine CG, which is a glycoprotein similar to equine pituitary LH, has been proven to have superior effects on follicle development in pigs. This gonadotrophin exhibits both LH- and FSH-like activities (*Farmer and Papkoff, 1979*) with a FSH/LH activity ratio that differs between 0.14 to 0.31 as estimated by bioassays conducted in rats and mice (*Bergfeld and Haring, 1983*). A different ratio of FSH versus LH activity may account for variations in the ovarian response seen in pigs after use of different eCG batches (*Bergfeld and Haring, 1983; Ciller et al., 2008*), and should be considered when analyzing the effectiveness of different treatment protocols that are based on eCG. As evident from numerous East-German experiments, 800 to 1,000 IU of eCG are most effective at stimulating follicle development in gilts and 600 to 1,000 IU eCG in sows. However, within- and between-farm differences occur, as indicated by variations in the ovarian response of sows that received the same eCG treatment but were located at different farms. This observation needs to be considered when determining an optimal dose of eCG (*Bergfeld et al., 1984*). It is recommended to test different eCG dosages first and then pick the one that yielded the optimum fertility. Besides the dose, the time when treatment occurs with eCG effects follicle development. For instance, the application of eCG one day after, or the last day of methallibure feeding in gilts (*Polge et al., 1968*) or 48 versus 24 hrs after methallibure (*Brüßow and Bergfeld, 1984*) was associated with an increased number of stimulated follicles and also with a lower dosages of eCG needed for

stimulation. Whenever a protocol includes the use of eCG it needs to be evaluated, for example by evaluating the ovary at slaughter, laparotomy, laparoscopy and ultrasound.

Besides pure eCG, different combinations of the gonadotrophins eCG and hCG have been used for stimulation of follicle development and estrus in gilts and sows (*Estienne et al., 2001; Knox et al., 2001*). However, using such combinations there is evidently the risk of inducing ovarian cysts and/or of premature luteinization of follicles. The latter is assumed due to the higher LH activity of those combinations compared to pure eCG, most likely as the result of the extra hCG (*Bergfeld et al., 1982*). Increasing eCG and hCG in a combination from 400 to 700 IU and from 200 to 350 IU, respectively, was associated with an increased number of gilts with cystic ovarian degeneration (36 % versus 88 %) and a decrease in the pregnancy rate (50 % versus 65 %). In contrast, when gilts were treated with 1,000 IU eCG, only 4 % of them developed ovarian cysts (*Schlegel et al., 1978*). Moreover, if 1,000 IU eCG was given to primi- and multiparous sows 24 hrs after weaning and compared to 400 IU eCG/200 IU hCG (Suigonan®, Intervet, Unterschleissheim, Germany), the pregnancy rates, litter sizes and piglet indices were all higher in sows that received eCG only (*Barbe et al., 1997*). Taken together the majority of the studies that were conducted in former East-Germany suggest that pure eCG stimulates follicle development with better fertility results than do combinations of eCG and hCG. This is certainly the reason why standardized, lyophilized eCG preparation (e.g. Pregmagon®; IDT Dessau, Germany) is given preference in the German pig industry for the purpose of stimulating follicle development in pigs. However, in the United States, a drug that combines 400 IU eCG and 200 IU hCG, i.e. PG 600 (Intervet Canada, Guelph, Ontario, Canada) is the preferred medication, and has been shown being effective for a multitude of reproductive purposes (*Kirkwood, 1999*).

Synthetically produced GnRH agonists have been used to stimulate follicle development in pigs, and are considered as a possible alternative to eCG. Recently, a synthetic analogue of lamprey GnRH-III (Peforelin; Maprelin® XP10, Veyx-Pharma, Schwarzenborn, Germany), was shown to exhibit selective FSH releasing activity in barrows (*Kauffold et al., 2005*). However, other studies in barrows (*Barretero-Hernandez et al., 2010*) and in gilts (*Brüssow et al., 2010*) did not provide any evidence of selective FSH release. Use of Maprelin® XP10 in primi- and multiparous sows to stimulate follicle development and estrus yielded in some experiments similar and in others opposing fertility results than eCG (*Engl et al., 2006; Kruse and Brüssow, unpublished*). However, a general recommendation for its broad use would greatly benefit from further supportive field results.

Induction of ovulation. Although, the onset of estrus and to some extent also ovulation can be synchronized in both gilts and sows by using eCG, the time when ovulation occurs can be still extremely variable. If fixed-time insemination is the goal, ovulation needs to be induced, either by using gonadotrophins with predominately LH activity such as hCG or by using GnRH

analogues. Human CG (*Hunter, 1967*) or equivalent substances such as pituitary extracts (*Tanabe et al., 1949*), aiming to mimic the endogenous preovulatory LH-peak, were effective in gilts at inducing ovulation which occurred at approximately 40-42 hrs after treatment. If animals are treated with eCG to stimulate follicle development and then followed by hCG treatment to induce ovulation, the interval between both treatments is however crucial to the ovulation-inducing effect of hCG. Considering that hCG should be given as close as possible to the time when the endogenous LH peak occurs in order to induce ovulation. Indeed, when this interval has been set at 78-80 hrs after eCG and the responses compared to non-treated controls, treated animals ovulated more uniformly, with ovulations occurring 42-53 hrs after hCG, whereas no control animal ovulate during this time period (*Bergfeld et al., 1976*).

Porcine LH (pLH, Lutropin-V, Bioniche Animal Health, Belleville, Canada) has been shown to be as effective as hCG at synchronizing ovulation in weaned sows (*Cassar et al., 2005; Bennett-Steward et al., 2008*). If eCG was given the day of weaning and pLH 80 hrs later, sows ovulated between 34-42 hrs post pLH (*Cassar et al., 2005*).

After GnRH, a decapeptide, was discovered and its structure known, the pharmaceutical industry started to synthesize GnRH and thus made it available for use in swine industry. In contrast to hCG, GnRH acts at the pituitary level and stimulates the release of endogenous LH, thereby approximating what is a more "biologically normal" event than does hCG. Once more the East-German pig industry took the initiative in the development of one of the first GnRH analogues, i.e. Gn-RH vet "Berlin-Chemie" for use in swine in the late 70s. *Brüßow and Bergfeld (1979)* and *Bergfeld and Brüßow (1979)* performed intensive studies on the effect of different doses of this GnRH analogue alone or in combination with hCG on ovulation and observed that, 900 µg of this GnRH analogue alone or different combinations of GnRH/hCG (100-300 µg GnRH/100-300 IU hCG) stimulate ovulation similar to what was achieved with only hCG.

Later, another GnRH agonist (D-Phe6-LHRH, Gonavet®, Berlin-Chemie, Berlin, Germany) was demonstrated to be even more effective at synchronizing ovulation in swine than Gn-RH vet "Berlin-Chemie". As observed by repeated laparoscopy (*Brüßow et al., 1990*), in gilts treated with 50 µg D Phe6-LHRH started ovulating 35.5 ± 2.7 hrs after treatment and finished ovulation on average 5.9 ± 1.7 hrs later. However, animals varied in their response to D-Phe6-LHRH, with variations being related to the interval between the GnRH injection and the LH peak, the maximum of the LH peak and the overall time needed for ovulation (Table 1). Field trials involving a total of 2,744 gilts that were all injected with 50 µg D-Phe6-LHRH (Gonavet®) 78-80 hrs after 1,000 IU eCG and artificially inseminated twice at fixed times, i.e. 24 and 40 hrs after GnRH, demonstrated that D-Phe6-LHRH yields superior fertility results than if ovulation was induced with hCG. A similar observation has been made with 71,600 sows that were treated with 50 µg D-Phe6-LHRH compared to 300 µg GnRH/ 300 IU hCG combination 55-58

hrs after eCG and artificially inseminated twice at 24 and 42 hrs after GnRH (Brüssow *et al.*, 1996). Other GnRH analogues that have been tested for ovulation induction in swine are busserelin (Möller-Holtkamp *et al.*, 1995; Martinat-Botte *et al.*, 2010), goserelin (Brüssow *et al.*, 2007) and triptorelin (Taibl *et al.*, 2008). All of them are effective at stimulating preovulatory LH secretion in both gilts and sows. Currently however, D-Phe6-LHRH is still the only GnRH analogue that has been licensed for use in swine in a number of European countries including Germany. The intravaginal application of GnRH containing gel (Baer and Bilkei, 2004; Taibl *et al.*, 2008) has been tested for the purpose of ovulation induction, but did not reach a stage beyond research.

Table 1. Effects of 50 µg D-Phe6-LHRH (Gonavet®) on LH and ovulation in gilts (from Brüssow *et al.*, 1994)

Criteria	Early* responder (n=6)	Medium* responder (n=6)	Late* responder (n=4)
GnRH – LH surge (hrs)	2.2 ± 0.4 ^a	5.0 ± 1.0 ^b	9.0 ± 1.0 ^c
LH maximum (ng/ml)	19.6 ± 9.8 ^a	7.3 ± 3.2 ^b	3.8 ± 0.8 ^c
GnRH – commencement of ovulation (hrs)	35.6 ± 2.4 ^a	34.6 ± 2.5 ^a	39.6 ± 1.5 ^b
GnRH – completion of ovulation (hrs)	39.0 ± 1.8 ^a	37.2 ± 2.5 ^a	42.0 ± 1.5 ^b
LH peak - commencement of ovulation (hrs)	33.4 ± 2.5	29.4 ± 3.9	30.6 ± 0.6
LH peak – completion of ovulation (hrs)	35.8 ± 1.7 ^a	32.2 ± 2.7 ^b	35.5 ± 3.8
Duration of ovulation (hrs)	3.6 ± 2.3	2.8 ± 1.7	2.4 ± 0.2

a,b p < 0.05,

* Animals were classified as early, medium and late responder according to their LH peak as response to GnRH.

Another neuronal peptide, kisspeptin, which is a product of the KISS1 gene (Kotani *et al.*, 2001) and synthesized as a preprohormone, has gained increasing interest lately since it is involved in the regulation of GnRH release, and if given to female pigs has been shown to stimulate LH release in a dose dependent manner (Lents *et al.*, 2008). Though kisspeptin is thus a promising candidate to induce ovulation in pigs the question of whether or not it will ever be used in the pig industry for this purpose as part of a fixed-time insemination protocol needs further investigation.

Protocols for fixed-time ovulation and insemination in practice

The German pig industry has a long and strong history of using biotechnology in pig reproduction. Supported by continuing research efforts, the industry has many years experience with this type of technology (Hühn *et al.*, 1996; Brüssow *et al.*, 1996), and has thus been able to recommend different protocols that can be used for estrus synchronization and fixed-time insemination

in pigs (Table 2). However, other approaches are also possible such as the one reported recently and performed with 3,000 sows on commercial farms in Canada (Cassar *et al.*, 2005; Bennett-Steward *et al.*, 2008). In these studies, sows were injected with 600 IU eCG at the day of weaning and were given 5 mg pLH 80 hrs later. They were then inseminated at fixed-times 36 and 44 hrs after pLH. Compared to untreated controls, the farrowing rate of treated sows was significantly increased (86 % versus 69 %). Similar high gestation rates (92 versus 96 %) were achieved when gilts were fixed-time inseminated 24 and 48 hrs after GnRH treatment with or without previous eCG stimulation (Martinat-Botte *et al.*, 2010).

Table 2. Treatment protocols for ovulation induction and fixed-time insemination in gilts and sows as recommended for use in practice based on experience from the East-German swine industry

Method	Gilts	Sows
Synchronisation of oestrous cycle	18 days oral application of altrenogest (20 mg/gilt/ day) (08:00 h)	Lactation until weaning
	15 days oral application of altrenogest (16 mg/gilt/day) (08:00 h)	
Stimulation of follicle development	800 – 1,000 IU eCG 24 hrs after last altrenogest (08:00 h)	Primiparous sows 1,000 IU eCG 24 hrs after weaning (08:00 h)
		Multiparous sows 600 – 800 IU eCG 24 hrs after weaning (08:00 h)
Induction of ovulation	GnRH* or hCG** 78-80 hrs after eCG (14:00-16:00 h)	Lactation > 4 weeks GnRH* or hCG** 56-58 hrs after eCG (16:00-18:00 h)
		Lactation 4 weeks GnRH* or hCG** 72 h after eCG (1600 h)***
		Lactation 3 weeks GnRH* or hCG** 78-80 hrs after eCG (14:00-16:00 h)
1st AI	24-26 hrs after GnRH* or hCG**	24-26 hrs after GnRH* or hCG**
2nd AI	38-40 hrs after GnRH* or hCG**	40-42 hrs after GnRH* or hCG**

* e.g. 50 µg Gonavet®, ** 500 IU hCG, *** weaning one day earlier in the afternoon

Conclusion

A technology is currently available that allows for manipulation of almost all key reproductive processes in the female pig including estrus and ovulation, making fixed-time insemination possible. Batch farrowing systems may profit from using this technology due, at least in part, to savings on labor and production of large batches of uniformly developed and healthy pigs. This technology does however not allow for compensation of health problems and/or mismanagement. Understanding of reproductive processes and the availability of hormonally active drugs are essential requirements when deciding to use this technology. However, several factors related to the farms themselves, as well as to the drugs may influence the effectiveness and outcome of this technology. It is thus recommended to adapt this technology to each individual farm.

Biološka i tehnološka pozadina sinhronizacije estrusa i fiksno vreme indukcije ovulacije u svinja

K.P. Brüßow, M. Wähler

Rezime

Tehnologija koja omogućava manipulisanje estrusom i ovulacijom, i koja takođe omogućava fiksno vreme osemenjavanja, može da bude od velike koristi za svinjarske farme koje raspolazu velikim brojem krmača. Takva tehnologija delimično štiti porođaj i omogućava proizvodnju velikog broja svinja istog uzrasta. Zahvaljujući dosadašnjim saznanjima o endokrinoj regulaciji razvoja folikula i ovulaciji, kao i dostupnim brojnim reproduktivno aktivnim supstancama, ova tehnologija je danas moguća. Ova "reproduktivna biotehnologija" će biti prikazana. Ona obuhvata postupke sinhronizacije estrusa zasnovane na upotrebi altrenogesta kod nazimica i velikog broja odlučениh krmača, za stimulaciju razvoja folikula koristi se eCG a za izazivanje ovulacije koristi se hCG ili LH kao i GnRH analog. Dok su postupci sinhronizacije estrusa pojedinačni, drugi postupci zahtevaju dodatni tretman. Ako je cilj fiksno vreme osemenjavanja, onda estrus treba da bude sinhronizovan a razvoj folikula i ovulacija da budu izazvani upotrebom GnRH analoga i hCG i da se ovulacija javlja nakon 36-42 sata. Opšta je preporuka da se osemenjavanje ovih životinja obavlja dvokratno, odnosno 24 i 40 časova nakon izazivanja ovulacije. Međutim, ova tehnologija zahteva zdrave životinje i jak menadžment i ne može se primeniti kao zamena za loš menadžment na farmi.

References

- BARBE C., WÄHNER M., SCHNURRBUSCH U. (1997): Influence of PMSG and PMSG/hCG-combinations on fertility of weaned sows. 2. Effect of different methods of cycle stimulation after weaning on fertility. *Arch. Tierzucht*, 40, 567-580.
- BAER C., BILKEI G. (2004): The effect of intravaginal applied GnRH-agonist on the time of ovulation and subsequent reproductive performance of weaned multiparous sows. *Reprod. Dom. Anim.*, 39, 293-297.
- BARRETERO-HERNANDEZ R., VIZCARRA J.A., BOWEN A., GALYEAN M. (2010): Lamprey GnRH-III releases luteinizing hormone but not follicle stimulating hormone in pigs. *Reprod. Dom. Anim.*, 45, 983-987.
- BENNET-STEWARD K., ARAMINI J., PELLAND C., FRIENDSHIP R. (2008): Equine chorionic gonadotrophin and porcine luteinizing hormone to shorten and synchronize the wean-to-breed interval among parity-one and paritytwo sows. *J. Swine Health Prod.*, 16, 182-187.
- BERGFELD J., FALGE R., HÜHN R., KAUFFOLD P. (1976): Experimental biological engineering to synchronise ovulation in gilts III Onset of ovulation, ovarian findings and fertilization following methallibure, 750 IU PMS with and without 500 IU hCG at different times of treatment. *Arch. Exper. Vet.-med.*, 30, 471-480.
- BERGFELD J., BRÜSSOW K.P. (1979): Preclinical studies into synchronisation of ovulation in gilts following treatment with Suisynchron PMSG and hCG/Gn-RH combination. *Mh. Vet. Med.*, 34, 613-615.
- BERGFELD J., BRÜSSOW K.P., TIEMANN U., RAASCH M.L. (1982): Orientation studies into selection of parameters for assessment of mixed gonadotrophic preparations for their biological effects on female fattening pigs. *Arch. Exper. Vet. Med.*, 36, 25-34.
- BERGFELD J., HARING E. (1983): Determination of gonadotrophic valence of preparations made of serum of pregnant mares (PMSG) III Biological determination of activity of "PMSG substandard Dessau" in bioassays for folliclestimulating hormone (FSH) and luteinizing hormone (LH). *Arch. Exper. Vet. Med.*, 37, 801-806.
- BERGFELD J., GEORGE G., BRÜSSOW K.P. (1984): Studies into dynamics and potency of ovulation in groups of sows on several farms following differentiated treatment to stimulate ovulation. *Arch. Exper. Vet. Med.*, 38, 849-856.
- BRÜSSOW K.P., BERGFELD J. (1979): Preliminary studies into Gn-RH and its use to release ovulation in prepuberal gilts and those with synchronised oestrus. *Mh. Vet.-Med.*, 34, 611-613.
- BRÜSSOW K.P., BERGFELD J. (1984): Studies into time shifting of ovulations in gilts by modification of injection dates in the context of synchronised ovulation. *Arch. Exper. Vet. Med.*, 38, 840-848.

- BRÜSSOW K.P., RÁTKY J., KANITZ E., BECKER F. (1990): The relationship between the surge of LH induced by exogenous Gn-RH and the duration of ovulation in gilts. *Reprod. Dom. Anim.*, 25, 255-260.
- BRÜSSOW K.P., KANITZ E., RÁTKY J. (1994): The dynamic of plasma luteinizing hormone surge and its relationships to the time of ovulation in gilts following exogenous GnRH. *Arch. Tierz.*, 37, 55- 63.
- BRÜSSOW K.P., JÖCHLE W., HÜHN U. (1996): Control of ovulation with a GnRH analog in gilts and sows. *Theriogenology*, 96, 925-934.
- BRÜSSOW K.P., SCHNEIDER F., NÜRNBERG G. (2001): Alteration of gonadotrophin and steroid hormone release, and of ovarian function by a GnRH antagonist in gilts. *Anim. Reprod. Sci.*, 66, 117-128.
- BRÜSSOW K.P., SCHNEIDER F., TUCHSCHERER A., RÁTKY J., KRAELING R.R., KANITZ W. (2007): Luteinizing hormone release after administration of the gonadotropinreleasing hormone agonist Fertilan (goserelin) for synchronization of ovulation in pigs. *J. Anim. Sci.*, 85, 129-137.
- BRÜSSOW K.P., SCHNEIDER F., TUCHSCHERER A., KANITZ W. (2010): Influence of synthetic Lamprey GnRH-III on gonadotropin release and steroid hormone levels in gilts. *Theriogenology*, 74, 1570-1578.
- CASSAR G., KIRKWOOD R.N., POLJAK Z., BENNETT-STEWARD K., FRIENDSHIP R.M. (2005): Effect of single or double insemination on fertility of sows bred at an induced estrus and ovulation. *J. Swine Health Prod.*, 13, 254-258.
- CILLER U.A., CILLER I.M., MCFARLANE J.R. (2008): Equine chorionic gonadotrophin isoform composition in commercial products compared with isoform composition in pregnant mare plasma. *Reprod. Fertil. Dev., Suppl.* 20, 77.
- DRIANCOURT M.A., LOCATELLI A., PRUNIER A. (1995): Effects of gonadotrophin deprivation on follicular growth in gilts. *Reprod. Nutr. Dev.*, 35, 663-673.
- ENGL S., KAUFFOLD J., SOBIRAJ A. ZAREMBA W. (2006): Investigations into the suitability of a GnRH-variant for estrus induction in pluriparous sows. *Reprod. Dom. Anim.*, 41, Suppl. 1, 33.
- ESTIENNE M.J., HARPER A.F., HORSLEY B.R., ESTIENNE C.E., KNIGHT J.W. (2001): Effects of PG600 on the onset of estrus and ovulation rate in gilts treated with Regumate. *J. Ani. Sci.*, 79, 2757-2761.
- ESTILL C.T. (1999): Current concepts in estrus synchronization in swine. *Proc Amer. Soc. Anim. Sci.*, 1-9.
- FARMER S.W., PAPKOFF H.H. (1979): Immunochemical studies with pregnant mare serum gonadotropins. *Biol. Reprod.*, 21, 425-431.
- GUTHRIE H.D., BOLT D.J. (1990): Changes in plasma follicle stimulating hormone, luteinizing hormone, estrogen and progesterone during growth of ovulatory follicles in the pig. *Dom. Anim. Endocrinol.*, 7, 83-91.

- HÜHN U., JÖCHLE W., BRÜSSOW K.P. (1996): Techniques developed for the control of estrus, ovulation and parturition in the East German pig industry: a review. *Theriogenology*, 46, 911-924.
- HUNTER R.H.F. (1967): The porcine ovulation after injection of human chorionic gonadotrophin. *Vet. Rec.*, 81, 21-23.
- HUNTER R.H.F. (1974): Chronological and cytological details of fertilization and early embryonic development in the domestic pig, *Sus scrofa*. *Anatom. Rec.*, 178, 169-186.
- KAUFFOLD J., SCHNEIDER F., ZAREMBA W., BRÜSSOW K.P. (2005): Lamprey GnRH-III stimulates FSH secretion in barrows. *Reprod. Dom. Anim.*, 40, 475-479.
- KIRKWOOD R.N. (1999): Pharmacological intervention in swine reproduction. *J. Swine Health Prod.*, 7, 29-35.
- KNOX R.V., RODRIGUEZ-ZAS S.L., MILLER G.M., WILLENBURG K.L., ROBB J.A. (2001): Administration of PG 600 to sows at weaning and the time of ovulation as determined by transrectal ultrasound. *J. Anim. Sci.*, 79, 796-802.
- KOTANI M., DETHEUX M., VANDENBOGAERDE A., COMMUNI D., VANDERWINDEN J.M., LE POUL E., BREZILLON S., TYLDESLEY R., SUAREZ-HUERTA N., VANDEPUT F., BLANPAIN C., SCHIFFMANN S.N., VASSART G., PARMENTIER M. (2001): The metastasis suppressor gene *KiSS-1* encodes kisspeptins, the natural ligands of the orphan G protein-coupled receptor *GPR54*. *J. Biol. Chem.*, 276, 34631-34636.
- LENTS C.L., HEIDORN N.L., BARB C.R., FORD J.J. (2008): Central and peripheral administration of kisspeptin activates gonadotropin but not somatotropin secretion in prepubertal gilts. *Reprod.*, 135, 879-887.
- LUCY M.C., LIU J., BOYD C.K., BRACKEN C.J. (2001): Ovarian follicular growth in sows. *Reprod.*, Suppl. 58, 31-45.
- MARTINAT-BOTTE F., BARITEAU F., FORGERIT Y., MACAR C., MOREAU A., TERQUI M., SIGNORET J.P. (1990): Oestrus control in gilts II synchronization of oestrus with the progestagen altrenogest (Regumate) effect on fertility and litter size. *Anim. Reprod. Sci.*, 22, 227-233.
- MARTINAT-BOTTÉ F., VENTURI E., GUILLOUET P., DRIANCOURT M.A., TERQUI M. (2009): Induction and synchronization of ovulations of nulliparous and multiparous sows with an injection of gonadotropin-releasing hormone agonist (Receptal). *Theriogenology*, 73, 332-342.
- MÖLLER-HOLTKAMP P., STICKAN F., PARVIZI N., ELSAESSER F. (1995): Release of LH in pigs after administration of the GnRH agonist buserelin in comparison to D-Phe6-LHRH. *Reprod. Dom. Anim.*, 30, 21-24.
- POLGE C., DAY B.N., GROVES T.W. (1968): Synchronisation of ovulation and artificial insemination in pigs. *Vet. Rec.*, 83, 136-142.
- POLGE C., DAY B.N. (1969): Induction of estrus and ovulation in swine during pituitary suppression with methallibure. *J. Anim. Sci.*, 69, 73-75.

-
- PRUNIER A., QUESNEL H. (2000): Nutritional influences on hormonal control of reproduction in female pigs. *Livest. Prod. Sci.*, 63, 1-16.
- SCHLEGEL W., WÄHNER M., STENZEL S. (1978): Different combinations of PMS and hCG – use for cycle stimulation in synchronized ovulation of gilts and effects on results of pregnancy. *Mh. Vet. Med.*, 33, 663-664.
- SCHWARTZ T., KOPYRA M., NOWICKI M. (2008): Physiological mechanisms of ovarian follicular growth in pigs – a review. *Acta Vet. Hung.*, 56, 369-378.
- TAIBL J.N., BREEN S.M., WEBEL S.K., KNOX R.V. (2008): Induction of ovulation using a GnRH agonist for use with fixed time AI in weaned sows. *Theriogenology*, 70, 1400.
- TANABE T.Y., WARNICK A.C., CASIDA L.E., GRUMMER R.H. (1949): The effect of gonadotrophins administered to sows and gilts during different stages of the estrual cycle. *J. Anim. Sci.*, 8, 550-557.
- VAN DEN BRAND H., DIELEMANN S.J., SOEDE N.M., KEMP B. (2000): Dietary energy source at two feeding levels during lactation in primiparous sows: I Effect on glucose, insulin, and luteinizing hormone and on follicle development, weaning-to-estrus interval, and ovulation rate. *J. Anim. Sci.*, 78, 396-404.
- VARLEY M.A., FOXCROFT G.R. (1990): Endocrinology of the lactating and weaned sow. *J. Reprod. Fertil., Suppl.* 40, 47-61.

Received 30 June 2011; accepted for publication 15 August 2011