

INFLUENCE OF REARING CONDITIONS ON REPRODUCTION, GROWTH, MILK YIELD AND QUALITY OF MEAT AND MILK OF SHEEP AND GOATS

Slavča Hristov¹, Branislav Stanković¹, Dimitar Nakov², Ivana Milošević-Stanković¹, Nevena Maksimović³

¹University of Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Belgrade, Republic of Serbia

²Ss.Cyril and Methodius University in Skopje, Faculty of Agricultural Sciences and Food, 1000 Skopje, Republic of Northern Macedonia

³Institute for Animal Husbandry, Autoput 16, 11080 Belgrade, Republic of Serbia

Corresponding author: Slavča Hristov, hristovs@agrif.bg.ac.rs

Review paper

Abstract: The most important stressors in our country are unfavourable climatic and nutritional conditions. Extreme deviations in climatic and nutritional conditions are immediate stressors for the body of sheep and goats. Appetite in sheep and goats decreases if they are exposed to extremely high ambient temperatures, while cold increases the body's energy needs and leads to the mobilization of fat from body fat and the consequent oxidation of fat and the formation of non-esterified fatty acids. Thyroid activity is reduced when these animals are exposed to high temperatures, and increased when they are exposed to cold. In an environment with high temperatures, blood glucose and cholesterol concentrations decrease due to unsuccessful maintenance of homeostasis. Sheep and goats are naturally kept in a group, which basically protects them from predators. They successfully register the action of new stimuli, especially noise and sudden movements of nearby predators. Early weaning breaks the bond between mother and offspring and adversely affects the development of normal behavioural relationships in lambs and kids, because artificial feeding systems do not fully meet the needs of oral activity, so sucking of the umbilical and scrotal region often occurs. Investigations of the influence of various technological procedures on the reactivity of the adrenal cortex indicate that the separation of the animal from the herd and immobilization are the most significant stimuli. Electrical stunning of sheep at the slaughterhouse causes significant stimulation of the sympathetic system with a sudden release of adrenaline and noradrenaline. In conditions of intensive production, the most significant stressors and physiological conditions that lead to stress are: parturition, birth, puerperal period, weaning, oestrus, high pregnancy, various surgical procedures, veterinary and zootechnical procedures, extreme variations of micro- and macroclimatic conditions, variations in quantity and food quality, etc.

Key words: rearing conditions, reproduction, production, sheep, goats

Introduction

In our climatic conditions, sheep and goats are raised on pastures for a longer period of the year, where the most significant stressors are unfavourable climatic and nutritional conditions (*Hristov and Bešlin, 1991*). Climate variations directly affect these species, influencing their biological reactions (*Hristov et al., 2002*). Detailed studies of certain haematological and biochemical changes related to pregnancy and lactation in sheep and goats in different climatic conditions, different rearing and housing conditions, as well as different feeding conditions have been performed. Periods requiring an abundant supply of nutrients, such as immediately before and after birth in highly productive breeds of sheep and goats under conditions of insufficient and/or unbalanced diet, may include mobilization of body fat and protein, leading to fatty liver and clinical ketosis (known as pregnancy toxæmia) and results in decreased productivity and in some cases even death (*Helman et al., 1995*). Similar metabolic changes can occur in low-yielding breeds of sheep and goats if these animal species are malnourished and/or fed unbalanced rations (*Hristov and Bešlin, 1991*).

Influence of stressors on reproduction, growth, milk yield and meat and milk quality

The connection between stress and animal welfare is a well-known issue in scientific observations (*Hristov et al., 2007*). Climate change is one of the most significant stressors affecting the welfare and production of sheep and goats (*Hristov et al., 2007a*). The impact of climate change is manifested through numerous losses in livestock production. It should be borne in mind that analyze of the conditions of production is an important issue, since each breed has different characteristics of adaptability. Changes in biological functions when exposed to heat stress include physiological, hormonal, haematological and biochemical reactions, which make sheep and goats resilient and enable them to survive in an unfavourable environment (*Darcan et al., 2007*). The extreme variations in climatic and nutritional conditions are the direct stress to the organism of sheep and goats (*Hristov and Bešlin, 1991*).

Appetite in sheep and goats, for example, decreases if they are exposed to extremely high ambient temperatures (*Sarangi, 2018*). Cold, on the other hand, increases the body's energy needs and leads to the mobilization of fat from body fat in sheep and goats and the consequent oxidation of fat, which leads to the formation of non-esterified fatty acids (NEFA – Non Esterified Fatty Acids) as a

constituent of blood plasma. NEFA level in blood plasma of sheep or goat is reliable indicator of nutritive status and reactivity to environmental stressors (Gupta *et al.*, 2013; Milošević-Stanković *et al.*, 2020).

The intensity of the detrimental effects of high environmental temperatures on sheep and goats depends on the efficiency of their thermoregulatory mechanisms (Caroprese *et al.*, 2009; Shaji *et al.*, 2017). Thyroid activity is reduced when these animals are exposed to high temperatures, and increased when they are exposed to cold (Habibu *et al.*, 2016; El-Tarabany *et al.*, 2017). Well-adapted animals react quickly to changes in the environment so they can make the necessary physiological adjustments. Research shows that in the environment with high temperatures, the concentrations of glucose and cholesterol in the blood decrease, which points out the unsuccessful maintenance of homeostasis.

Heat stress has a greater impact on total cholesterol concentration, which may be due to increased use of fatty acids for energy production and decreased glucose concentration in animals exposed to heat stress (Gupta *et al.*, 2013; Das *et al.*, 2016). Similarly, researchers point out protein level increase when these animals are exposed to heat stress (Nazifi *et al.*, 2003; Gupta *et al.*, 2013). During physical exertion a faster release of oxygen is present as well as an increase of oxygen consumption and thus an increase of the concentration of haemoglobin. Reduced food intake in animals under long-term heat stress reduces erythrocyte production and haemoglobin levels, resulting in a reduction in the number of red blood cells in the bloodstream (Alam *et al.*, 2011; Gupta *et al.*, 2013).

Lambs and kids born by caesarean section or exposed to cold have an increased concentration of NEFA in the blood plasma (Clarke *et al.*, 1994; Hristov *et al.*, 2012). Other adverse factors such as, for example, complete lack of food or malnutrition combined with multiple pregnancy also lead to a significant increase in the concentration of NEFA in the blood plasma of sheep and goats, which is closely related to the intensity of stressors (Eşki *et al.*, 2015; Di Giacomo *et al.*, 2016; Milošević Stanković *et al.*, 2020). In sheep and goats exposed to cold as well as humid climate conditions, the adrenal cortex is stimulated to increase hormone secretion while the number of eosinophils in the blood is significantly reduced. The metabolic reaction to repeated exposures of sheep and goats to cold in a short duration is progressively reduced. Progressive changes in climate conditions are likely to lead to relatively wide ranges in haematological values in sheep and goats in different geographic areas (Ghosh *et al.*, 2014; Snyman *et al.*, 2017).

Increased cortisol levels in sheep and goats occur after acute heat stress (Marai *et al.*, 2007; Seijan *et al.*, 2008); intensive chronic stress can result in periods of high concentrations of cortisol, followed by a decrease in individual health condition through immuno-suppression and atrophy of the body's defence tissues. In addition, cortisol reduces the energy available to the immune system, which increases the susceptibility of animals to infectious diseases.

Today, the most intensive production in sheep and goat breeding has contributed to the production of lambs and kids throughout the year. Basically, the existing systems in use in the housing and rearing of sheep and goats during the year and the artificial induction of oestrus have contributed to the reduction of exposure of sheep and goats to daylight (*Alcalde et al., 2017*).

Highly productive breeds of sheep and goats have been subjected to rigorous selection and crossbreeding with tested rams and goats to improve meat quality. The lambs and kids are weaned and fed with artificial milk using rubber teats from the second day of age. A study of the behaviour of groups of lambs and kids reveals that housing systems where the social bond between mother and offspring is broken, adversely affect the development of normal behavioural relationships (*Sevi et al., 2009*). As it was found for calves (*Johnsen et al., 2016*), artificial feeding systems do not fully meet the needs of oral activity of lambs and kids, since it was noticed that those lambs and kids develop the habit of sucking the umbilical and scrotal regions by individual lambs and kids in the group (*Stephens and Baldwin, 1971; Napolitano et al., 2008*). Previous attempts to improve the diet in order to achieve the most optimal in accordance with the biological needs of sheep and goats have not been completely successful in preventing the occurrence or reduction of abnormal oral activity of lambs and kids (*Miranda-de la Lama and Mattiello, 2010*).

Surgical transplantation of the adrenal gland to the neck where the blood supply is easy enabled the use of sheep as a model for studying adrenocortical function in mammals. The relatively simple surgical procedure made it possible for the uterus to connect with the adrenal cortex of the foetus and consequently easily stimulate with ACTH and induce a reaction in the foetus and newborn lambs (*McDonald et al., 1958*). Studies of the dynamics of adrenal gland reactions to experimental ACTH stimulation in adult sheep showed that the maximum secretion of cortisol is achieved within 10 to 20 minutes. When the ACTH infusion is continuous, maximum cortisol secretion is maintained for the next 30 to 60 minutes, after which normal secretion is slowly established (*Beaven et al., 1964; Salemi et al., 2000*).

Sheep and goats are naturally kept in a group, which basically protects them from predators. They have a very developed observation about the action of new stimuli, especially noise and sudden movements of nearby predators (*Rutter, 2002*). Sudden loud noise, such as a loud gunshot, is a stressor and causes a significant initial reaction in sheep movement. Depending on the duration and intensity of the sound, the herd moves quickly in different durations and at different distances (*Hargreaves and Hutson, 1990; Grandin, 2011*).

Aberrations from stationary level limits of many constituents in blood plasma have also been found in sheep and goats exposed to qualitatively new stimuli. An increase in thyroid hormone release rate was found in sheep within 15 to 30 minutes, when they were exposed to a series of fireworks explosions or barking dogs. The increase in the concentration of these hormones usually lasts for about 2

hours (Falconer and Hetzel, 1964; Bobek et al., 1986). Plasma NEFA concentrations also increase in sheep and goats when exposed to qualitatively new stimuli, such as vein puncture (Paterson, 1963). This increase usually does not last longer than 2 hours (Slee and Halliday, 1968), and the increase rate becomes progressively lower with repeated seizures in order to take blood samples (Reid and Hinks, 1962; Westgren et al., 1987). Similarly, the consequent progressive decrease was caused by venous puncture when taking blood to determine the concentration of corticosteroids in the blood plasma (Bassett and Hinks, 1969).

Circadian variations in plasma cortisol concentrations in sheep and goats are absent for the first 2 to 3 days after the introduction of noise and sudden movements of nearby predators. During the stay in the new ambient conditions, the diurnal rhythm of cortisol concentration in the blood plasma is reestablished. However, a significantly higher concentration of cortisol in blood plasma is found in unadapted individuals compared to the concentration in fully adapted sheep and goats. It is considered that in sheep and goats require a period of 7 to 28 days for complete adaptation to the new conditions of keeping and accommodation (McNatty and Young, 1973).

In the study of various technological procedures and their effects on the reactivity of the adrenal cortex, it was found that the separation of sheep and goats from the herd and immobilization are the most influential stimuli. Prolonged shear time from 5 to 15 minutes causes an increase in plasma cortisol concentration by 50% (Kilgour and de Langen, 1970).

It has been proven that an increase in the concentration of cortisol in the blood plasma of sheep can be induced by an intravenous infusion of adrenaline. The increasing the concentration of cortisol was dependent on the applied dose of adrenaline, as well as higher in sheep that were housed in the stall. Studies have shown that clinical signs of disorders and disturbances in sheep are partly related to the release of adrenaline. The researchers point out that significant dynamic adrenocortical activity is found when a group of sheep is exposed to disturbance and suggest that the adrenal gland response depends on the intensity of the stressor (Thurley and McNatty, 1973).

The use of inexperienced dogs to chase the flock in sheep causes a greater and longer-lasting response of the organism than the use of trained dogs. Grouping of sheep by humans causes a less stressful reaction of the sheep organism (Beausoleil et al., 2005). Surprisingly, sheep kept together with lambs, although more often in a distressed state, show less reaction to grouping with experienced dogs compared to the same treatment in the same sheep before parturition.

Sheep housed in stables immediately before transport are less susceptible to stress or increased corticosteroid concentrations. However, sheep caught and loaded into a trailer directly on pasture show a greater reaction (Miranda-de la Lama et al., 2012; Pascual-Alonso et al., 2017).

Electrical stunning of sheep at the slaughterhouse causes significant stimulation of the sympathetic system with a sudden release of adrenaline and noradrenaline into the circulation, 14 to 20 times higher than normal values. Concentrations of cortisol, adrenaline and noradrenaline in blood plasma are significantly higher in sheep in large slaughterhouses compared to the values found in the same species in small non-specialized slaughterhouses. This is thought to be due to heavier catching, higher noise and greater disturbance of sheep in large slaughterhouses (*Linares et al., 2008; Sabow et al., 2016*).

Mean concentrations of adrenal hormones in the blood serum of sheep gradually increase with an increase in the duration of transport to 2 hours. When extending the duration of transport to 8 hours, a lower rate of increase in the concentration of these hormones in the blood serum of sheep was observed (*Shannon et al., 1976; Zarogiannis et al., 2006; Kadim et al., 2007*).

Meat breeds of sheep and goats are inevitably exposed to various procedures before and during loading, during transport, during unloading, as well as food deprivation before slaughter. Procedures with sheep and goats before, during and after transport can affect animal welfare and production profitability. Expressed stress before slaughter negatively affects the quality of meat in animals. Weight loss during transport and storage in pre-slaughter depots are of particular importance because in small ruminants the digestive tract represents a larger proportion of live weight than in cattle and pigs (*Ekiz and Yalcintan, 2013; Hashem et al., 2013*).

The researchers used several physiological indicators to assess stress levels in farm animals. Cortisol concentration was used as a reliable indicator of short-term physical stress (*Kannan et al., 2003*). Leptin, a protein hormone synthesized by adipose tissue, has been reported to inhibit the activity of the hypothalamic-pituitary-adrenal axis in response to stress. (*Macedo et al., 2019*). Creatine kinase (CK), an enzyme in skeletal muscle, is released into the blood plasma in response to muscle damage under stress (*Nazifi et al., 2003; Chulayo and Muchenje, 2013*). Stress in sheep and goats due to lack of food causes the degradation of proteins and increase the level of urea in the blood (*McDougal et al., 1991; Wheelock et al., 2010*). The effects of various stressors on leukocyte profiles in farm animals were determined. Physiological responses to pre-slaughter stress have been studied in all species of farm animals, with limited data on goats (*Ekiz and Yalcintan, 2013; Sabow et al., 2016*).

Immediately before and after birth, highly productive breeds of sheep and goats need large amounts of nutrients. If then there is insufficient and/or unbalanced diet, body fat and protein are mobilized, which leads to fatty liver and clinical ketosis (known as pregnancy toxemia), reduced productivity, and in some cases the death of these species (*Brozos et al., 2011; Milošević-Stanković et al., 2020*). Similar metabolic changes can occur in poorly productive breeds of sheep and goats if they are not fed and/or fed an unbalanced diet. Concentrations of free

fatty acids, triglycerides, cholesterol, urea and creatinine in the blood serum are higher in sheep and goats with the occurrence of pregnancy ketosis and multiple pregnancies compared to non-pregnant sheep and goats (*Donoghue and Kronfeld, 1990; Brozos et al., 2011*). Calcium intake decreases with reduced food intake during ketosis, which leads to hypocalcaemia. Serum calcium levels were found to be significantly reduced in sheep and goats with hypocalcaemia and pregnancy toxemia. Also, reduced food intake during grazing is a possible cause of hypocalcaemia (*Kessler, 1991*).

Conclusion

Based on a detailed review of the literature data related to the influence of sheep and goat rearing conditions as a stressor on reproduction, growth, milk yield and quality of meat and milk, the following can be concluded:

- Extreme deviations in climatic and nutritional conditions are immediate stressors for sheep and goats;
- Appetite in sheep and goats is reduced if they are exposed to extremely high ambient temperatures. Cold, on the other hand, increases the body's energy needs and leads to the mobilization of fat from body reserve in sheep and goats and the consequent oxidation of fat, which causes the formation of non-esterified fatty acids;
- Thyroid activity is reduced when these animals are exposed to high temperatures and increased when they are exposed to cold. In an environment with high temperatures, blood glucose and cholesterol concentrations decrease and homeostasis fails;
- Sheep and goats have very good registration of the action of new stimuli, especially noise and sudden movements of nearby predators;
- Breaking the social bond between mother and offspring by early weaning adversely affects the development of normal behavioural relationships in lambs and kids;
- Separation of sheep and goats from the herd and immobilization are the most influential stress stimuli;
- Electrical stunning of sheep and goats at the slaughterhouse causes significant stimulation of the sympathetic system with a sudden release of adrenaline and noradrenaline into the circulation.

In the conditions of modern intensive sheep and goat production, the most significant stressors, regardless of origin, and physiological conditions in which these species are more susceptible to stress are: parturition, birth, puerperal period, weaning, oestrus, high pregnancy, various surgical procedures, veterinary and zootechnical procedures, extreme variations of micro- and macroclimatic conditions, variations in the quantity and quality of food, etc.

Uticaj uslova gajenja ovaca i koza na reprodukciju, prirast, mlečnost i kvalitet mesa i mleka

Slavča Hristov, Branislav Stanković, Dimitar Nakov, Ivana Milošević-Stanković, Nevena Maksimović

Rezime

Najvažniji stresori kod nas su nepovoljni klimatski i nutritivni uslovi. Ekstremna odstupanja u klimatskim i nutritivnim uslovima predstavljaju neposredne stresore za organizam ovaca i koza. Apetit kod ovaca i koza se smanjuje ako su one izložene izrazito visokim ambijentalnim temperaturama, dok hladnoća povećava energetske potrebe organizma i dovodi do mobilizacije masti iz masnog tkiva organizma i posledične oksidacije masti i stvaranja neesterifikovanih masnih kiselina. Aktivnost štitne žlezde je smanjena kada su ove životinje izložene visokim temperaturama, a povećana kada su izložene hladnoći. U ambijentu sa visokim temperaturama smanjuju se koncentracije glukoze i holesterola u krvi usled neuspešnog održavanja homeostaze. Ovcе i koze se prirodno drže u grupi čime se u osnovi štite od predatora. One uspešno registruju delovanje novih stimulusa, naročito buke i iznenadnih pokreta predatora u blizini. Rano odbijanje prekida vezu majke i mladunčeta i nepovoljno utiče na razvoj normalnih bihejvioralnih odnosa kod mladunčadi, jer veštački sistemi ishrane ne zadovoljavaju u potpunosti potrebe oralne aktivnosti, pa se često javlja sisanje pupčane i skrotalne regije. Ispitivanja uticaja različitih tehnoloških postupaka na reaktivnost kore nadbubrežne žlezde ukazuju da je izdvajanje jedinke iz stada i imobilizacija predstavljaju najznačajnije stimulse. Električno omamljivanje ovaca na klanici izaziva znatnu stimulaciju simpatičkog sistema sa naglim oslobađanjem adrenalina i noradrenalina. U uslovima intenzivne proizvodnje najznačajniji stresori i fiziološka stanja koja dovode do stresa su: porodaj, rođenje, puerperalni period, odbijanje mladunčadi, estrus, visoki graviditet, razni hirurški postupci, veterinarske i zootehničke mere, ekstremna variranja mikro- i makroklimatskih uslova, variranja u količini i kvalitetu hrane i dr.

Ključne reči: uslovi držanja, reprodukcija, proizvodnja, ovce, koze

Acknowledgement

This paper was supported by the Serbian Ministry of Education and Science, grant TR31086.

References

- ALAM M.M., HASHEM M.A., RAHMAN M.M., HOSSAIN M.M., HAQUE M.R., SOBHAN Z., ISLAM M. S. (2011): Effect of heat stress on behaviour, physiological and blood parameters of goat. *Progressive Agriculture*, 22, 1-2, 37-45.
- ALCALDE M.J., SUÁREZ M.D., RODERO E., ÁLVAREZ R., SÁEZ M.I., MARTÍNEZ T.F. (2017): Effects of farm management practices and transport duration on stress response and meat quality traits of suckling goat kids. *Animal*, 11, 1626-1635.
- BASSETT J.M., HINKS N.T. (1969): Micro-determination of corticosteroids in ovine peripheral plasma: effects of venipuncture, corticotropin, insulin and glucose. *Journal of Endocrinology*, 44, 387.
- BEAUSOLEIL N.J., STAFFORD K.J., MELLOR D.J. (2005): Sheep show more aversion to a dog than to a human in an arena test. *Applied Animal Behaviour Science*, 91, 3-4, 219-232.
- BEAVEN D.W., ESPINER E.A., HART D.S. (1964): The suppression of cortisol secretion by steroids, and response to corticotrophin in sheep with adrenal transplants. *The Journal of Physiology*, 171, 2, 216.
- BOBEK S., NIEZGODA J., PIERZCHAŁA K., LITYNSKI P., SECHMAN A. (1986): Changes in circulating levels of iodothyronines, cortisol and endogenous thiocyanate in sheep during emotional stress caused by isolation of the animals from the flock. *Journal of Veterinary Medicine Series A*, 33, 1-10, 698-705.
- BROZOS C., MAVROGIANNI V.S., FTHENAKIS G.C. (2011): Treatment and control of peri-parturient metabolic diseases: pregnancy toxemia, hypocalcemia, hypomagnesemia. *Veterinary Clinics: Food Animal Practice*, 27, 1, 105-113.
- CAROPRESE M., CASAMASSIMA D., PIER GIACOMO RASSU S., NAPOLITANO F., SEVI A. (2009): Monitoring the on-farm welfare of sheep and goats. *Italian Journal of Animal Science*, 8(sup1), 343-354.
- CHULAYO A.Y., MUCHENJE V. (2013): The effects of pre-slaughter stress and season on the activity of plasma creatine kinase and mutton quality from different sheep breeds slaughtered at a smallholder abattoir. *Asian-Australasian Journal of Animal Sciences*, 26, 12, 1762.

- CLARKE L., DARBY C.J., LOMAX M.A., SYMONDS M.E. (1994): Effect of ambient temperature during 1st day of life on thermoregulation in lambs delivered by cesarean section. *Journal of Applied Physiology*, 76, 4, 1481-1488.
- DARCAN N., CEDDEN F., GUNEY O. (2007): Spraying effects on goat welfare in hot and humid climate. *American Journal of Animal and Veterinary Sciences*, 2, 4, 99-103.
- DAS R., SAILO L., VERMA N., BHARTI P., SAIKIA J. (2016): Impact of heat stress on health and performance of dairy animals: A review. *Veterinary World*, 9, 3, 260.
- DI GIACOMO K., SIMPSON S., LEURY B.J., DUNSHEA F.R. (2016): Dietary betaine impacts the physiological responses to moderate heat conditions in a dose dependent manner in sheep. *Animals*, 6, 9, 51.
- DONOGHUE S., KRONFELD D.S. (1990): Clinical nutrition of sheep and goats. *Veterinary Clinics of North America: Food Animal Practice*, 6, 3, 563-576.
- EKIZ E. E., YALCINTAN H. (2013): Comparison of certain haematological and biochemical parameters regarding pre-slaughter stress in saanen, maltese, gokceada and hair goat kids. *Istanbul Üniversitesi Veteriner Fakültesi Dergisi*, 39, 189-196.
- EL-TARABANY M.S., EL-TARABANY A.A., ATTA M.A. (2017): Physiological and lactation responses of Egyptian dairy Baladi goats to natural thermal stress under subtropical environmental conditions. *International Journal of Biometeorology*, 61, 1, 61-68.
- EŞKI F., TAŞAL İ., KARSLI M.A., ŞENDAĞ S., USLU B.A., WAGNER H., WEHREND A. (2015): Concentrations of NEFA, β -HBA, triglycerides, and certain blood metabolites in healthy colored Angora goats during the peripartum period. *Turkish Journal of Veterinary and Animal Sciences*, 39, 4, 401-405.
- FALCONER I.R., HETZEL B.S. (1964): Effect of emotional stress and TSH on thyroid vein hormone level in sheep with exteriorized thyroids. *Endocrinology*, 75, 42-48.
- GHOSH S., SINGH A.K., HALDAR C. (2014): Seasonal modulation of immunity by melatonin and gonadal steroids in a short day breeder goat *Capra hircus*. *Theriogenology*, 82, 8, 1121-1130.
- GRANDIN T. (2011): Understanding flight zone and point of balance for low stress handling of cattle, sheep, and pigs. Online <http://www.grandin.com> (Retrieved: 17 May 2014).
- GUPTA M., KUMAR S., DANGI S.S., JANGIR B.L. (2013): Physiological, biochemical and molecular responses to thermal stress in goats. *International Journal of Livestock Research*, 3, 27-38.
- HABIBU B., KAWU M.U., MAKUN H.J., ALUWONG T., YAQUB L. S. (2016): Seasonal variation in body mass index, cardinal physiological variables and serum thyroid hormones profiles in relation to susceptibility to thermal stress in goat kids. *Small Ruminant Research*, 145, 20-27.

- HARGREAVES A.L., HUTSON G.D. (1990): The stress response in sheep during routine handling procedures. *Applied Animal Behaviour Science*, 26, 1-2, 83-90.
- HASHEM M.A., HOSSAIN M.M., RANA M.S., ISLAM M.S., SAHA N.G. (2013): Effect of heat stress on blood parameter, carcass and meat quality of Black Bengal goat. *Bangladesh Journal of Animal Science*, 42, 1, 57-61.
- HELMAN R.G., ADAMS L.G., BRIDGES C.H. (1995): The lesions of hepatic fatty cirrhosis in sheep. *Veterinary Pathology*, 32, 6, 635-640.
- HRISTOV S. (2002): Stres ovaca. *Zoohigijena. Univerzitet u Beogradu, Poljoprivredni fakultet, Zemun-Beograd.*, 368-370.
- HRISTOV S., BEŠLIN R. (1991): Stres domaćih životinja. *Monografija. Poljoprivredni fakultet. Zemun-Beograd*, 157.
- HRISTOV S., MAKSIMOVIĆ N., STANKOVIĆ B., ŽUJOVIĆ M., PANTELIĆ V., STANIŠIĆ N., ZLATANOVIĆ Z. (2012): The most significant stressors in intensive sheep production. *Biotechnology in Animal Husbandry*, 28, 4, 649-658.
- HRISTOV S., FRATRIĆ N., KIROVSKI D., STANKOVIĆ B. (2007): Stres i dobrobit farmskih životinja. *Tematski zbornik "Dobrobit životinja i biosigurnost na farmama"*, Prva međunarodna konferencija o dobrobiti i biosigurnosti na farmama u Srbiji, Poljoprivredni fakultet, Beograd, 87-95.
- HRISTOV S, VUČINIĆ M, STANKOVIĆ B. (2007a): Zašto nam je potrebna dobrobit životinja. *Tematski zbornik "Dobrobit životinja i biosigurnost na farmama"*, Prva međunarodna konferencija o dobrobiti i biosigurnosti na farmama u Srbiji, Poljoprivredni fakultet, Beograd, 5-21
- JOHNSEN J.F., ZIPP K.A., KÄLBER T., DE PASSILLÉ A.M., KNIERIM U., BARTH K., MEJDELL C.M. (2016): Is rearing calves with the dam a feasible option for dairy farms?—Current and future research. *Applied Animal Behaviour Science*, 181, 1-11.
- KADIM I.T., MAHGOUB O., ALKINDI A.Y., AL-MARZOOQI W., AL-SAQRI N.M., ALMANEY M., MAHMOUD I.Y. (2007): Effect of transportation at high ambient temperatures on physiological responses, carcass and meat quality characteristics in two age groups of Omani sheep. *Asian-Australasian Journal of Animal Sciences*, 20, 3, 424-431.
- KANNAN G., KOUAKOU B., TERRILL T.H., GELAYE S. (2003): Endocrine, blood metabolite, and meat quality changes in goats as influenced by short-term, preslaughter stress. *Journal of Animal Science*, 81, 6, 1499-1507.
- KESSLER J. (1991): Mineral nutrition of goats. *Goat nutrition*, 46, 104-119.
- KILGOUR R., DE LANGEN H. (1970): Stress in sheep resulting from management practices. *Proceedings of the New Zealand Society of Animal Production* 30, 65-76.
- LINARES M.B., BÓRNEZ R., VERGARA H. (2008): Cortisol and catecholamine levels in lambs: Effects of slaughter weight and type of stunning. *Livestock Science*, 115, 1, 53-61.

- MACEDO T.J.S., SANTOS J.M.S., BEZERRA M.É.S., MENEZES V.G., GOUVEIA B.B., BARBOSA L.M.R., BARBOSA G.L.T.L, MONTEA A.P.O., BARBERINO R.S., BATISTAB A.M., BARROSA V.R.P., WISCHRALB A., QUEIROZC M.A.A., ARAÚJOD G.G.L., MATOS M.H.T. (2019): Immunolocalization of leptin and its receptor in the sheep ovary and in vitro effect of leptin on follicular development and oocyte maturation. *Molecular and cellular endocrinology*, 495, 110506.
- MARAI I.F.M., EL-DARAWANY A.A., FADIEL A., ABDEL-HAFEZ M.A.M. (2007): Physiological traits as affected by heat stress in sheep—a review. *Small ruminant research*, 71, 1-3), 1-12.
- MCDONALD I.R., GODING J.R., WRIGHT R.D. (1958): Transplantation of the adrenal gland of the sheep to provide access to its blood supply. *Australian Journal of Experimental Biology and Medical Science*, 36, 1, 83-96.
- MCDUGALL S., LEPHERD E.E., SMITH S. (1991): Haematological and biochemical reference values for grazing Saanen goats. *Australian Veterinary Journal*, 68, 11, 370-372.
- MCNATTY K.P., YOUNG A. (1973): Diurnal changes of plasma cortisol levels in sheep adapting to a new environment. *Journal of Endocrinology*, 56, 329-330.
- MILOŠEVIĆ-STANKOVIĆ I., HRISTOV S., MAKSIMOVIĆ N., POPOVIĆ B., DAVIDOVIĆ V., MEKIĆ C., DIMITRIJEVIĆ B., CINCOVIĆ M., STANKOVIĆ B. (2020): Energy metabolism indicators and body condition in periparturient period of Alpine goats. *Large Animal Review*, 26, 1, 13-18.
- MIRANDA-DE LA LAMA G.C., MATTIELLO S. (2010): The importance of social behaviour for goat welfare in livestock farming. *Small Ruminant Research*, 90, 1-3, 1-10.
- MIRANDA-DE LA LAMA G.C., SALAZAR-SOTELO M.I., PÉREZ-LINARES C., FIGUEROA-SAAVEDRA F., VILLARROEL M., SAÑUDO C., María G.A. (2012): Effects of two transport systems on lamb welfare and meat quality. *Meat Science*, 92, 4, 554-561.
- NAPOLITANO F., DE ROSA G., SEVI A. (2008): Welfare implications of artificial rearing and early weaning in sheep. *Applied Animal Behaviour Science*, 110, 1-2, 58-72.
- NAZIFI S., SAEB M., ROWGHANI E., KAVEH K. (2003): The influences of thermal stress on serum biochemical parameters of Iranian fat-tailed sheep and their correlation with triiodothyronine (T3), thyroxine (T4) and cortisol concentrations. *Comparative Clinical Pathology*, 12, 3, 135-139.
- PASCUAL-ALONSO M., MIRANDA-DE LA LAMA G.C., AGUAYO-ULLOA L., VILLARROEL M., MITCHELL, M., MARÍA G.A. (2017): Thermophysiological, haematological, biochemical and behavioural stress responses of sheep transported on road. *Journal of Animal Physiology and Animal Nutrition*, 101, 3, 541-551.

- PATTERSON D.S.P. (1963): Some observations on the estimation of nonesterified fatty acid concentration in cow and sheep plasma. *Research in Veterinary Science*, 4, 230-237.
- REID R.L., HINKS N.T. (1962): Studies on the carbohydrate metabolism of sheep. XVII Feed requirements and voluntary feed intake in late pregnancy, with particular reference to prevention of hypoglycaemia and hyperketonaemia. *Australian Journal of Agricultural Research*, 13, 1092-1111.
- RUTTER S.M. (2002): *Behaviour of Sheep and Goats 10. The ethology of domestic animals: An introductory text*, 145.
- SABOW A.B., GOH Y.M., ZULKIFLI I., SAZILI A.Q., KAKA U., AB KADI M.Z.A., EBRAHIMI M., NAKYINSIGEI K., ADEYEMI K.D. (2016): Blood parameters and electroencephalographic responses of goats to slaughter without stunning. *Meat science*, 121, 148-155.
- SALEMI R., MCDUGALL J.G., HARDY K.J., WINTOUR E.M. (2000): Effect of adrenomedullin infusion on basal and stimulated aldosterone secretion in conscious sheep with cervical adrenal autotransplants. *Journal of Endocrinology*, 166, 2, 389-400.
- SARANGI S. (2018): Adaptability of goats to heat stress: A review. *Pharma Innovation*, 7, 4, 1114-1126.
- SEJIAN V., SRIVASTAVA R.S., VARSHNEY V.P. (2008): Pineal-adrenal relationship: modulating effects of glucocorticoids on pineal function to ameliorate thermal-stress in goats. *Asian-Australasian Journal of Animal Sciences*, 21, 7, 988-994.
- SEVI A., CASAMASSIMA D., PULINA G., PAZZONA A. (2009): Factors of welfare reduction in dairy sheep and goats. *Italian Journal of Animal Science*, 8 (sup1), 81-101.
- SHAJI S., SEJIAN V., BAGATH M., MANJUNATHAREDDY G.B., KURIEN E.K., VARMA G., BHATTA R. (2017): Summer season related heat and nutritional stresses on the adaptive capability of goats based on blood biochemical response and hepatic HSP70 gene expression. *Biological Rhythm Research*, 48, 1, 65-83.
- SHANNON A.D., QUIN J.W., JONES M.A.S. (1976): Response of the regional lymphatic system of the sheep to acute stress and adrenaline. *Quarterly Journal of Experimental Physiology and Cognate Medical Sciences: Translation and Integration*, 61, 3, 169-183.
- SLEE J., HALLIDAY R. (1968): Some effects of cold exposure, nutrition and experimental handling on serum free fatty-acid levels in sheep. *Animal Production*, 10, 1, 67-76.
- SNYMAN M.A., STORBECK K., SWART P. (2017): Physiological parameters of three South African Angora goat CYP17 genotypes subjected to cold stress. *Grootfontein Agricultural*, 17, 1, 1-10.

STEPHENS D.B., BALDWIN B.A. (1971): Observations on the Behaviour of Groups of Artificially Reared. Lambs. *Research in Veterinary Science*, 12, 3, 219-225.

THURLEY D.C., MCNATTY K.P. (1973): Factors affecting peripheral cortisol levels in unrestricted ewes. *Acta Endocrinology*, (Copenh), 74, 2, 331-7.

WESTGREN M., ALMSTROM H., NYMAN M., ULMSTEN U. (1987): Maternal perception of sound-provoked fetal movements as a measure of fetal well-being. *BJOG: An International Journal of Obstetrics & Gynaecology*, 94, 6, 523-527.

WHEELLOCK J.B., RHOADS R.P., VANABLE M.J., SANDERS S.R., BAUGARD L.H. (2010): Effects of heat stress on energetic metabolism in lactating Holstein cows. *Journal of Dairy Science*, 93, 2, 644-655.

ZAROGIANNIS S., HATZOGLOU C., STEFANIDIS I., MATAFIA G., VOGIATZIDIS K., GOURGOULIANIS K., MOLYVDAS P.A. (2006): Effect of adrenaline on transmesothelial resistance of isolated sheep pleura. *Respiratory Physiology & Neurobiology*, 150, 2-3, 165-172.

Received 9 December 2020; accepted for publication 20 December 2020